


Dietary Reference Intakes for Calcium and Vitamin D



Calcium and vitamin D are two essential nutrients long known for their role in bone health. Over the last ten years, the public has heard conflicting messages about other benefits of these nutrients—especially vitamin D—and also about how much calcium and vitamin D they need to be healthy.

To help clarify this issue, the U. S. and Canadian governments asked the Institute of Medicine (IOM) to assess the current data on health outcomes associated with calcium and vitamin D. The IOM tasked a committee of experts with reviewing the evidence, as well as updating the nutrient reference values, known as Dietary Reference Intakes (DRIs). These values are used widely by government agencies, for example, in setting standards for school meals or specifying the nutrition label on foods. Over time, they have come to be used by health professionals to counsel individuals about dietary intake.

The committee provided an exhaustive review of studies on potential health outcomes and found that the evidence supported a role for these nutrients in bone health but not in other health conditions. Overall, the committee concludes that the majority of Americans and Canadians are receiving adequate amounts of both calcium and vitamin D. Further, there is emerging evidence that too much of these nutrients may be harmful.

Overall, the committee concludes that the majority of Americans and Canadians are receiving adequate amounts of both calcium and vitamin D. Further, there is emerging evidence that too much of these nutrients may be harmful.

Health Effects of Vitamin D and Calcium Intake

The new reference values are based on much more information and higher-quality studies than were available when the values for these nutrients were first set in 1997. The committee assessed more than one thousand studies and reports and listened to testimony from scientists and stakeholders before

making its conclusions. It reviewed a range of health outcomes, including but not limited to cancer, cardiovascular disease and hypertension, diabetes and metabolic syndrome, falls, immune response, neuropsychological functioning, physical performance, preeclampsia, and reproductive outcomes. This thorough review found that information about the health benefits beyond bone health—benefits often reported in the media—were from studies that provided often mixed and inconclusive results and could not be considered reliable. However, a strong body of evidence from rigorous testing substantiates the importance of vitamin D and calcium in promoting bone growth and maintenance.

Dietary Reference Intakes

The DRIs are intended to serve as a guide for good nutrition and provide the basis for the development of nutrient guidelines in both the United States and Canada. The science indicates that on average 500 milligrams of calcium per day meets the requirements of children ages 1 through 3, and on average 800 milligrams daily is appropriate for those ages 4 through 8 (see table for the Recommended Dietary Allowance—a value that meets the needs of most people). Adolescents need higher levels to support bone growth: 1,300 milligrams per day meets the needs of practically all adolescents. Women ages 19 through 50 and men up to 71 require on average 800 milligrams daily. Women over 50 and both men and women 71 and

TABLE: Dietary Reference Intakes for Calcium and Vitamin D

Life Stage Group	Calcium			Vitamin D		
	Estimated Average Requirement (mg/day)	Recommended Dietary Allowance (mg/day)	Upper Level Intake (mg/day)	Estimated Average Requirement (IU/day)	Recommended Dietary Allowance (IU/day)	Upper Level Intake (IU/day)
Infants 0 to 6 months	*	*	1,000	**	**	1,000
Infants 6 to 12 months	*	*	1,500	**	**	1,500
1-3 years old	500	700	2,500	400	600	2,500
4-8 years old	800	1,000	2,500	400	600	3,000
9-13 years old	1,100	1,300	3,000	400	600	4,000
14-18 years old	1,100	1,300	3,000	400	600	4,000
19-30 years old	800	1,000	2,500	400	600	4,000
31-50 years old	800	1,000	2,500	400	600	4,000
51-70 year old males	800	1,000	2,000	400	600	4,000
51-70 year old females	1,000	1,200	2,000	400	600	4,000
>70 years old	1,000	1,200	2,000	400	800	4,000
14-18 years old, pregnant/lactating	1,100	1,300	3,000	400	600	4,000
19-50 years old, pregnant/lactating	800	1,000	2,500	400	600	4,000

*For infants, Adequate Intake is 200 mg/day for 0 to 6 months of age and 260 mg/day for 6 to 12 months of age.

**For infants, Adequate Intake is 400 IU/day for 0 to 6 months of age and 400 IU/day for 6 to 12 months of age.

This thorough review found that information about the health benefits beyond bone health—benefits often reported in the media—were from studies that provided often conflicting results and could not be considered reliable.

older should take in 1,000 milligrams per day on average to ensure they are meeting their daily needs for strong, healthy bones.

Determining intake levels for vitamin D is somewhat more complicated. Vitamin D levels in the body may come from not only vitamin D in the diet but also from synthesis in the skin through sunlight exposure. The amount of sun exposure one receives varies greatly from person to person, and people are advised against sun exposure to reduce the risk of skin cancer. Therefore, the committee assumed minimal sun exposure when establishing the DRIs for vitamin D, and it determined that North Americans need on average 400 International Units (IUs) of vitamin D per day (see table for the Recommended Dietary Allowances—values sufficient to meet the needs of virtually all persons). People age 71 and older may require as much as 800 IUs per day because of potential changes in people’s bodies as they age.

Questions About Current Intake

National surveys in both the United States and Canada indicate that most people receive enough calcium, with the exception of girls ages 9-18, who often do not take in enough calcium. In contrast, postmenopausal women taking supplements may be getting too much calcium, thereby increasing their risk for kidney stones.

Information from national surveys shows vitamin D presents a complicated picture. While the average total intake of vitamin D is below the median requirement, national surveys show that

average blood levels of vitamin D are above the 20 nanograms per milliliter that the IOM committee found to be the level that is needed for good bone health for practically all individuals. These seemingly inconsistent data suggest that sun exposure currently contributes meaningful amounts of vitamin D to North Americans and indicates that a majority of the population is meeting its needs for vitamin D. Nonetheless, some subgroups—particularly those who are older and living in institutions or who have dark skin pigmentation—may be at increased risk for getting too little vitamin D.

Before a few years ago, tests for vitamin D were conducted infrequently. In recent years, these tests have become more widely used, and confusion has grown among the public about how much vitamin D is necessary. Further, the measurements, or cut-points, of sufficiency and deficiency used by laboratories to report results have not been set based on rigorous scientific studies, and no central authority has determined which cut-points to use. A single individual might be deemed deficient or sufficient, depending on the laboratory where the blood is tested. The number of people with vitamin D deficiency in North America may be overestimated because many laboratories appear to be using cut-points that are much higher than the committee suggests is appropriate.

Tolerable Upper Levels of Intake

The upper level intakes set by the committee for both calcium and vitamin D represent the safe



Committee to Review Dietary Reference Intakes for Vitamin D and Calcium

A. Catharine Ross (Chair)
Professor of Nutrition and
Occupant of Dorothy Foehr
Huck Chair in Nutrition, Penn-
sylvania State University

Steven A. Abrams
Professor of Pediatrics, Baylor
College of Medicine

John F. Aloia
Professor, SUNY at Stony
Brook, Chief Academic Officer,
Winthrop-University Hospital

Patsy M. Brannon
Professor, Division of
Nutritional Sciences, Cornell
University

Steven K. Clinton
Professor, Division of Hematol-
ogy and Oncology, The Ohio
State University

Ramon A. Durazo-Arvizu
Associate Professor, Loyola
University Stritch School of
Medicine

J. Christopher Gallagher
Professor of Medicine, Creigh-
ton University Medical Center

Richard L. Gallo
Professor of Medicine and
Pediatrics, University of
California-San Diego

Glennville Jones
Head, Department of Biochem-
istry and Professor of Bio-
chemistry & Medicine, Queens
University, Ontario

Christopher S. Kovacs
Professor of Medicine (Endocri-
nology), Memorial University of
Newfoundland

Joann E. Manson
Professor of Medicine and the
Elizabeth Brigham Professor of
Women's Health, Harvard
Medical School

Susan T. Mayne
Professor of Epidemiology and
Public Health, Yale School of
Public Health

Clifford J. Rosen
Senior Scientist, Maine Medical
Center Research Institute

Sue A. Shapses
Professor, Department of
Nutritional Sciences, Rutgers
University

Consultant

Hector F. Deluca
University of Wisconsin-
Madison

Study Staff

Christine L. Taylor
Study Director

Ann L. Yaktine
Senior Program Officer

Heather B. DelValle
Associate Program Officer

Heather Breiner
Program Associate

Anton Bandy
Financial Officer

Geraldine Kennedo
Administrative Assistant, Food
and Nutrition Board

Linda D. Meyers
Director, Food and Nutrition
Board

Study Sponsors

Agricultural Research Service, U.S. Department of Agriculture
Center for Nutrition Policy and Promotion, U.S. Department of
Agriculture

Department of the Army, U.S. Department of Defense

Food and Drug Administration, U.S. Department of Health and
Human Services

Health Canada

Office of Disease Prevention and Health Promotion, U.S.
Department of Health and Human Services

National Institutes of Health (Division of Nutrition Research
Coordination, National Institute on Aging, National Institute of
Arthritis and Musculoskeletal and Skin, National Cancer
Institute, and Office of Dietary Supplements), U.S. Department of
Health and Human Services

boundary at the high end of the scale and should not be misunderstood as amounts people need or should strive to consume. While these values vary somewhat by age, as shown in the table, the committee concludes that once intakes of vitamin D surpass 4,000 IUs per day, the risk for harm begins to increase. Once intakes surpass 2,000 milligrams per day for calcium, the risk for harm also increases.

As North Americans take more supplements and eat more of foods that have been fortified with vitamin D and calcium, it becomes more likely that people consume high amounts of these nutrients. Kidney stones have been associated with taking too much calcium from dietary supplements. Very high levels of vitamin D (above 10,000 IUs per day) are known to cause kidney and tissue damage. Strong evidence about possible risks for daily vitamin D at lower levels of intake is limited, but some preliminary studies offer tentative signals about adverse health effects.

Conclusion

Scientific evidence indicates that calcium and vitamin D play key roles in bone health. The current evidence, however, does not support other benefits for vitamin D or calcium intake. More targeted research should continue. However, the committee emphasizes that, with a few exceptions, all North Americans are receiving enough calcium and vitamin D. Higher levels have not been shown to confer greater benefits, and in fact, they have been linked to other health problems, challenging the concept that “more is better.”

INSTITUTE OF MEDICINE

OF THE NATIONAL ACADEMIES

Advising the nation/Improving health

500 Fifth Street, NW
Washington, DC 20001

TEL 202.334.2352

FAX 202.334.1412

www.iom.edu

The Institute of Medicine serves as adviser to the nation to improve health.

Established in 1970 under the charter of the National Academy of Sciences, the Institute of Medicine provides independent, objective, evidence-based advice to policy makers, health professionals, the private sector, and the public.

Copyright 2011 by the National Academy of Sciences. All rights reserved.